

## Research Article

### Effects of Consumption of Cinnamon on Blood Glucose and Lipid profile of the Patients of Type 2 Diabetes

\* Gullapalli H. S. MD,<sup>1</sup> Avinash P. Tekade MD<sup>2</sup>, Namrata H. Gullapalli M.Sc.<sup>3</sup>

<sup>1</sup>Associate Professor, Department of Physiology, Vinayaka Mission's Medical college and Hospital, Karaikal, Puducherry.

<sup>2</sup>Associate Professor, Department of Physiology, Katuri Medical College and Hospital, Guntur, Andhra Pradesh.

<sup>3</sup>Dietitian and nutritionist Vinayaka Mission's Medical college and Hospital, Karaikal, Puducherry.

#### Corresponding author

Gullapalli H. S

Email: [hgullapalli2000@gmail.com](mailto:hgullapalli2000@gmail.com)

---

**Abstract – Objective**— The objective of this study was to determine whether cinnamon improves blood glucose, triglyceride, total cholesterol, HDL cholesterol, and LDL cholesterol levels in people with type 2 diabetes. **Research Design And Methods**—A total of 60 people with type 2 diabetes, 30 men and 30 women aged  $52.2 \pm 6.32$  years, were divided randomly into six groups. Groups 1, 2, and 3 consumed 1, 3, or 6 g of cinnamon daily, respectively, and groups 4, 5, and 6 were given placebo capsules corresponding to the number of capsules consumed for the three levels of cinnamon. The cinnamon was consumed for 40 days followed by a 20-day washout period. **Results**—After 40 days, all three levels of cinnamon reduced the mean fasting serum glucose (18–29%), triglyceride (23–30%), LDL cholesterol (7–27%), and total cholesterol (12–26%) levels; no significant changes were noted in the placebo groups. Changes in HDL cholesterol were not significant. **Conclusions**—The results of this study demonstrate that intake of 1, 3, or 6 g of cinnamon per day reduces serum glucose, triglyceride, LDL cholesterol, and total cholesterol in people with type 2 diabetes and suggest that the inclusion of cinnamon in the diet of people with type 2 diabetes will reduce risk factors associated with diabetes and cardiovascular diseases.

**Keywords** – Cinnamon, blood glucose, LDL-cholesterol, Insulin, Anti-diabetic activity

---

#### INTRODUCTION

It is postulated that spices may play a role in control of blood sugar level. Spices such as cinnamon, cloves, bay leaves, and turmeric display insulin-enhancing activity in vitro [12,3]. Aqueous extracts from cinnamon have been shown to increase in vitro glucose uptake and glycogen synthesis and to increase phosphorylation of the insulin receptor; in addition, these cinnamon extracts are likely to aid in triggering the insulin cascade system [11,12]. Because insulin also plays a key role in lipid metabolism, we postulated that consumption of cinnamon would lead to improved glucose and blood lipids in vivo. Therefore, this study was designed to determine whether there is a dose response of cinnamon on clinical variables associated with diabetes and cardiovascular diseases in people with type-2 diabetes. [1-10]

#### RESEARCH DESIGN AND METHODS

This study was conducted in the Department of Physiology, Vinayaka Mission Institute of Medical Sciences, Karaikal, Puducherry and was approved by the Ethics Committee. Selection criteria for the study included the following for people with type 2 diabetes: age >40 years, not on insulin therapy, not taking medicine for other health conditions, and fasting blood glucose levels between 7.8 and 22.2 mmol/l (140–400

mg/dl). A total of 60 individuals with type 2 diabetes, 30 men and 30 women, were selected for the study. The mean age of the subjects was  $52.0 \pm 6.87$  years in the placebo groups and  $52.0 \pm 5.85$  years in the groups consuming cinnamon. The duration of diabetes was also similar:  $6.73 \pm 2.32$  years for the placebo group and  $7.10 \pm 3.29$  years for the cinnamon groups. There were also an equal number of men and women in the placebo and cinnamon groups. All subjects were taking sulfonylurea drugs, i.e., glibenclamide; medications did not change during the study. Cinnamon (*Cinnamomum cassia*) certified by Spice Board, Kochi, Kerala. Cinnamon and wheat flour were ground finely and put into. Each capsule contained either 500 mg of cinnamon or wheat flour. Both the cinnamon and placebo capsules were packaged in plastic bags containing 40 capsules (1 g or two capsules per day for 20 days), 120 capsules (3 g or six capsules per day for 20 days), or 240 capsules (6 g or 12 capsules per day for 20 days) and prepared for distribution to the subjects. When subjects finished testing after the first 20 days, they were given the second package of capsules. Compliance was monitored by capsule count and contact with the subjects. Compliance was considered excellent and all capsules were consumed.

The study was conducted for 60 days with 60 individuals with type 2 diabetes divided randomly into

six equal groups. Group 1 consumed two 500-mg capsules of cinnamon per day, group 2 consumed six capsules of cinnamon per day, and group 3 consumed 12 capsules of cinnamon per day. Groups 4, 5, and 6 were assigned to respective placebo groups, which consumed a corresponding number of capsules containing wheat flour. Subjects consumed their normal diets and continued their medications throughout the study. From days 41 to 60, no cinnamon or placebo was given. The 1-g dose of cinnamon and placebo was spread over the day as 0.5 g (one capsule) after lunch and 0.5 g after dinner. The 3-g and 6-g doses of cinnamon and placebo were spread over the day as 1 g (two capsules) and 2 g (four capsules) after breakfast, lunch, and dinner, respectively. The subjects were instructed to take the capsules immediately after meals on days 0, 20, 40, and 60; ~5 ml of fasting blood was collected from each subject. Blood samples were transferred to sterilized centrifuge tubes and allowed to clot at room temperature. The blood samples were centrifuged for 10 min in a tabletop clinical centrifuge at 4,000 rpm for serum separation. Serum samples were stored in a freezer at 0°C for later analyses.

Glucose level was determined using an autoanalyzer (Express Plus; Ciba Corning Diagnostics, Palo Alto, CA). Triglyceride levels were determined by the enzymatic colorimetric method of Werner *et al.*[13]

using an autoanalyzer (Express Plus; Ciba Corning) and an Elitech kit (Meditek Instrument, India). Cholesterol levels were determined by enzymatic colorimetric method of Allain *et al*[14] using the same autoanalyzer. Chylomicrons, VLDL, and LDL were precipitated by adding phosphotungstic acid and magnesium ions to the sample. Centrifugation left only the HDL in the supernatant[15]. LDL cholesterol was calculated by dividing the triglycerides by 5 and subtracting the HDL cholesterol[16].

Two-way ANOVA and randomized complete block design were used for statistical analysis [17]. Values are means  $\pm$  SD.

## RESULTS

The addition of 1, 3, or 6 g of cinnamon to the diet led to significant decreases in serum glucose levels after 40 days. Values after 20 days were significantly lower only in the group receiving 6 g of cinnamon ([Table 1](#)). At the levels tested, there was no evidence of a dose response because the response to all three levels of cinnamon was similar. Decreases ranged from 18 to 29%. After the subjects no longer consumed the cinnamon for 20 days, glucose levels were significantly lower only in the group consuming the lowest level of cinnamon. Glucose values in the three placebo groups were not significantly different at any of the time points.

**Table -1: Effects of cinnamon on glucose levels in people with type 2 diabetes**

Group*	Dose of Cinnamon g/day	Fasting Blood Glucose Level (mmol/l)			
		Before Cinnamon Intake	During Cinnamon Intake		After Cinnamon Intake
		Day 0	Day 20	Day 40	Day 60
1	1	11.6 $\pm$ 1.7	10.5 $\pm$ 1.8	08.7 $\pm$ 1.6	09.7 $\pm$ 1.4
2	3	11.4 $\pm$ 1.2	09.9 $\pm$ 1.1	09.4 $\pm$ 1.1	09.9 $\pm$ 1.6
3	6	13.0 $\pm$ 1.4	10.1 $\pm$ 1.3	09.2 $\pm$ 1.5	11.4 $\pm$ 1.8
4	Placebo 1	12.2 $\pm$ 1.0	12.7 $\pm$ 0.8	12.4 $\pm$ 1.1	12.6 $\pm$ 1.0
5	Placebo 2	12.4 $\pm$ 1.0	11.8 $\pm$ 0.9	12.7 $\pm$ 1.0	12.6 $\pm$ 1.3
6	Placebo 3	16.7 $\pm$ 1.4	16.7 $\pm$ 1.6	16.8 $\pm$ 1.7	17.0 $\pm$ 1.3

Data are means  $\pm$  SD \* Ten individual in each group

The consumption of cinnamon also led to a time-dependent decrease in serum triglyceride levels at all amounts of cinnamon tested after 40 days ([Table 2](#)). Values after 20 days were significantly lower only in the group consuming 6 g of cinnamon per day. Decreases after 40 days of cinnamon consumption ranged from 23 to 30%. These data indicate that consumption of cinnamon for >20 days was more beneficial than shorter use for reduction of triglyceride

levels in people with type 2 diabetes. The mean fasting serum triglyceride levels of the subjects who consumed 1 g or 3 g of cinnamon per day for 40 days followed by 20 days of not consuming cinnamon were still significantly lower than the mean fasting serum triglyceride levels of the same groups at the beginning of the study. Decreases in the 6-g group were no longer significant. There were no changes in triglyceride levels in any of the three placebo groups ([Table 2](#)).

**Table -2: Effects of cinnamon on triglyceride levels in people with type 2 diabetes**

Group*	Doses of Cinnamon (g/day)	Fasting Serum Triglyceride Level (mmol/l)			
		Before Cinnamon Intake	During Cinnamon Intake		After Cinnamon Intake
		Day 0	Day 20	Day 40	Day 60
1	1	2.25±0.35	1.92±0.18	1.57±0.21	1.57±0.21
2	3	2.27±0.30	2.74±0.49	2.01±0.36	2.15±0.52
3	6	2.48±0.39	1.81±0.28	1.91±0.30	2.07±0.32
4	Placebo 1	2.31±0.32	2.38±0.34	2.50±0.30	2.45±0.32
5	Placebo 2	2.38±0.29	2.42±0.31	2.39±0.28	2.21±0.29
6	Placebo 3	2.25±0.34	2.66±0.38	2.52±0.40	2.55±0.35

Data are means ± SD \* Ten individual in each group

There were also significant decreases in serum cholesterol levels in all three groups consuming cinnamon, and no changes were noted in the respective placebo groups (Table 3). Decreases were significant after 20 days, and values were similar after 40 days,

except in the group consuming 3 g per day, which continued to decrease. These decreases in serum cholesterol level ranging from 13 to 26% were maintained even after not consuming additional cinnamon for 20 days (Table 3, last column).

**Table -3: Effects of cinnamon on cholesterol levels in people with type 2 diabetes**

Group*	Doses of Cinnamon (g/day)	Fasting Serum cholesterol Level (mmol/ml)			
		Before Cinnamon Intake	During Cinnamon Intake		After Cinnamon Intake
		Day 0	Day 20	Day 40	Day 60
1	1	4.91±0.23	4.32±0.21	4.32±0.27	4.09±0.30
2	3	5.51±0.29	4.76±0.32	4.09±0.26	4.03±0.34
3	6	5.30±0.22	4.63±0.21	4.55±0.24	4.86±0.19
4	Placebo 1	4.58±0.28	4.67±0.35	4.58±0.31	4.78±0.31
5	Placebo 2	4.81±0.30	4.71±0.30	5.04±0.31	4.94±0.35
6	Placebo 3	5.51±0.41	5.59±0.44	5.55±0.40	5.84±0.42

Data are means ± SD \* Ten individual in each group

Decreases in LDL were significant in the 3- and 6-g groups after 40 days with decreases of 10 and 24% (Table 4). Decreases in the 1-g group were not significant after 40 days but continued to decline during the washout period and were significant after 60 days

(Table 4, last column). There were non-significant changes in HDL in the subjects consuming 1 or 6 g of cinnamon for 40 days. Decreases in the 3-g group were significant after 20 days. These values remained relatively unchanged after the 20-day washout period.

**Table -4: Effects of cinnamon on LDL levels in people with type 2 diabetes**

Group*	Doses of Cinnamon (g/day)	Fasting Serum LDL Level (mmol/ml)			
		Before Cinnamon Intake	During Cinnamon Intake		After Cinnamon Intake
		Day 0	Day 20	Day 40	Day 60
1	1	2.55±0.12	2.28±0.15	2.48±0.10	2.35±0.13
2	3	2.77±0.18	2.43±0.28	2.04±0.19	1.97±0.18
3	6	2.87±0.18	2.55±0.13	2.59±0.15	2.72±0.11
4	Placebo 1	2.30±0.22	2.30±0.31	2.20±0.22	2.40±0.22
5	Placebo 2	2.55±0.25	2.40±0.22	2.55±0.27	2.79±0.27
6	Placebo 3	3.03±0.31	3.15±0.33	3.28±0.34	3.35±0.37

Data are means ± SD \* Ten individual in each group

## DISCUSSION:

This study demonstrates effects of low levels (1–6 g per day) of cinnamon on the reduction of glucose, triglyceride, LDL cholesterol, and total cholesterol

levels in subjects with type 2 diabetes. The study design serves to replicate the results because there were similar effects at the three doses tested. It is not clear whether even less than 1 g of cinnamon per day would also be beneficial. The data are also reinforced by the

observation that there were no significant changes in any of the placebo groups. There were also no problems with compliance or problems associated with the consumption of  $\leq 6$  g of cinnamon per day. The mechanism of the effects of cinnamon on glucose and blood lipids must be determined. Symptoms of insulin resistance include decreased stimulation of muscle glycogen synthesis as well as defects in glycogen synthase activity and glucose uptake<sup>18</sup>. In addition, altered enzymatic activities, such as an increased phosphatase activity and/or seryl phosphorylation of the insulin receptor substrate by glycogen synthase kinase-3 (GSK-3), have also been shown to be involved in some cases of type 2 diabetes [19,20]. Dephosphorylation of the receptor  $\beta$ -subunit is associated with the deactivation of its kinase activity and, therefore, is associated with insulin signal down regulation [21]. Maximal phosphorylation of the insulin receptor is associated with increased insulin sensitivity, which is associated with improved glucose and lipid levels. Extracts of cinnamon activated glycogen synthase, increased glucose uptake, and inhibited glycogen synthase kinase-3 $\beta$  [11,12]. Extracts of cinnamon also activated insulin receptor kinase and inhibited dephosphorylation of the insulin receptor, leading to maximal phosphorylation of the insulin receptor [12]. All of these effects would lead to increased insulin sensitivity. We have shown that extracts of cinnamon also function as potent antioxidants, which would lead to additional health benefits of this substance (unpublished data). Dhuley [22] showed that cinnamon displays antioxidant activity in rats fed a high-fat diet. The maintenance of lower serum glucose and lipid levels, even when the individuals were not consuming cinnamon for 20 days, denotes sustained effects of cinnamon, indicating that cinnamon would not need to be consumed every day. The levels of cinnamon tested in this study, 1–6 g per day, suggest that there is a wide range of cinnamon intake that may be beneficial and that intake of  $<1$  g daily is likely to be beneficial in controlling blood glucose and lipid levels.

**Conclusion:** cinnamon reduced serum glucose, triglyceride, total cholesterol, and LDL cholesterol levels in people with type 2 diabetes. Because cinnamon would not contribute to caloric intake, those who have type 2 diabetes or those who have elevated glucose, triglyceride, LDL cholesterol, or total cholesterol levels may benefit from the regular inclusion of cinnamon in their daily diet. In addition, cinnamon may be beneficial for the remainder of the population to prevent and control elevated glucose and blood lipid levels.

#### REFERENCES:

1. Raza A, Movahed A: Current concepts of cardiovascular diseases in diabetes mellitus. *Int J Cardiol* 89:123–134, 2003
2. Khan A, Bryden NA, Polansky MM, Anderson RA: Insulin potentiating factor and chromium

- content of selected foods and spices. *Bio Trace Element Res* 24:183–188, 1990
3. Broadhurst CL, Polansky MM, Anderson RA: Insulin-like biological activity of culinary and medicinal plant aqueous extracts in vitro. *J Agric Food Chem* 48:849–852, 2000 [CrossRefMedline](#)
4. Bailey CJ, Day C: Traditional plant medicines as treatments for diabetes. *Diabetes Care* 12:553–564, 1989 Shapiro K, Gong WC: Natural products used for diabetes. *J Am Pharm Assoc* 42:217–226, 2002 .
5. Shapiro K, Gong WC: Natural products used for diabetes. *J Am Pharm Assoc* 42:217–226, 2002
6. Rashwan AA: Effects of dietary additions of anise, fenugreek and caraway on reproductive and productive performance of New Zealand White rabbit does. *Egypt J Rabbit Sci* 8:157–167, 1998
7. Khan BA, Abraham A, Leelamma S: Biochemical response in rats to the addition of curry leaf (*Murraya koenigii*) and mustard seeds (*Brassica juncea*) to the diet. *Plant Foods Hum Nutr* 49:295–299, 1996
8. Khan BA, Abraham A, Leelamma S: Influence of spices—*Murraya koenigii* and *Brassica juncea*—on rats fed atherogenic diet. *J Food Sci* 35:66–68, 1998
9. Chithra V, Leelamma S: Hypolipidemic effect of coriander seeds (*Coriandrum sativum*): mechanism of action. *Plant Foods Hum Nutr* 51:167–172, 1997
10. Zhang WL, Liu DW, Wo XD, Zhang YH, Jin MM, Ding ZS: Effects of *Curcuma longa* on proliferation of cultured bovine smooth muscle cells and on expression of low-density lipoprotein receptor in cells. *Chinese Med J* 112:308–311, 1999
11. Imparl-Radosevich J, Deas S, Polansky MM, Baedke DA, Ingebrutsen TS, Anderson RA, Graves DJ: Regulation of phosphorylase phosphatase (PTP-1) and insulin receptor kinase by fractions from cinnamon: implications for cinnamon regulation of insulin signaling. *Horm Res* 50:177–182, 1998
12. Jarvill-Taylor KJ, Anderson RA, Graves DJ: A hydroxychalcone derived from cinnamon functions as a mimetic for insulin in 3T3-L1 adipocytes. *J Am Coll Nutr* 20:327–336, 2001
13. Werner M, Gabrielson DG, Eastman G: Ultramicrodeterminations of serum triglycerides by bioluminescent assay. *Clin Chem* 21:268–271, 1981
14. Allain CC, Poon LS, Chon CSG, Richmond U, Fu PC: Enzymatic determination of total serum cholesterol. *Clin Chem* 20:470–475, 1974
15. Lopes-Virella MF, Stone P, Ellis S, Coldwell JA: Cholesterol determinations in high density

- lipoproteins separated by three methods. Clin Chem 23:882–884, 1977
16. Friedewald WT, Levy RI, Fredrickson DS: Estimation of the concentration of low density lipoprotein cholesterol in plasma without the use of the preparative ultracentrifuge. Clin Chem 18:499–502, 1972
  17. Freed RD: MSTAT-C With MGRAPH. Version 2.00. East Lansing, MI, Michigan State University, 1997
  18. Cline GW, Oetersen KF, Krssak M, Shen J, Hundal RS, Trajanoski Z, Inzucchi S, Dresner A, Rothman DL, Shulman GI: Impaired glucose transport as a cause of decreased insulin-stimulated muscle glycogen synthesis in type 2 diabetes. N Engl J Med 341:240–245,
  19. Begum N, Sussman KE, Draznin B: Differential effects of diabetes on adipocyte and liver phosphotyrosine and phosphoserine phosphatase activities. Diabetes 40:1620–1629, 1991
  20. Nadiv O, Shinitzke M, Manu H, Hecht D, Roberts CT, LeRoith D, Zick Y: Elevated protein tyrosine phosphatase activity and increased membrane viscosity are associated with impaired activation of the insulin receptor kinase in old rats. Biochem J 298:443–450, 1994
  21. Eldar-Finkelman H, Krebs EG: Phosphorylation of insulin receptor substrate-1 by glycogen synthase kinase 3 impairs insulin action. Proc Natl Acad Sci 94:9660–9664, 1997
  22. Dhuley JN: Antioxidant effects of cinnamon (Cinnamomum verum) bark and greater cardamom (Amomum sabulatum) seeds in rats fed high fat diet. Indian J Exp Biol 37:238–242